



جامعة جنوب الوادي - كلية العلوم ببنها - قسم الجيولوجيا

مقرر الجيولوجيا التركيبية

لطلاب الفرقة الثالثة - قسم الجيولوجيا [لائحة جديدة]

جميع الشعب

الفصل الدراسي الأول

2024-2023

Geology Department

Structural Geology Course

for 3rd Year Students

by

Prof. Dr. Gamal M. Kamal El-Din

CONTENTS

Chapter- I Introduction.....	4
- Structural geology	
- What is structural geology?	
- Structure Vs Tectonics	
- Goals of structural geology	
- Importance of structural geology	
- Applications of Structural Geology	
- STRESS Vs Strain	
Chapter- II	
Tectonics.....	15
- Continental Drift Theory	
- Seafloor Spreading	
- Plate Tectonics	
Plate Boundaries	
Hotspots	
Tectonic setting of the Red Sea	
East African Rift System	
Chapter- III - Primary Structures.....	41
- Depositional Structures	
- Deformational Structures	
- Sedimentary Structures	
- Igneous Structures	
- Depositional Contacts	
Chapter- IV - Faults.....	52
- Introduction	
- BRITTLE deformation	
- DUCTILE deformation	
- Fault geometry	

Structural Geology Course for 3rd Year Students - Geology Department

- The Main Types of Faults
- Classification of Faults
- Strike – slip faults
 - San Andreas Fault
 - Dead Sea Fault
 - Najd Fault System
 - Pull-apart basins
- Reverse faults
 - Thrust faults
 - Zagros thrust belt
- Normal faults
 - Horst Vs Graben
 - Listric faults
- Stress distribution during faulting
- FAULT CRITERIA
 - Shear Zones

Chapter- V Folds.....96

- Introduction
- Geometry of a Fold (Parts of a Fold)
- Classifications of Folds
- Mechanism of Folding

Chapter- VI Foliation and lineation.....115

- Introduction
- Foliation Mechanism
- Lineation
- Types of lineation

References.....122

Chapter- I

Introduction

- Objectives

The aims of Structural Geology are the study of deformed rocks. This is achieved by the description of the geometry of geological structures, by the kinematics that rocks have experienced during their deformation history and by the understanding of the dynamics involved during the deformation. There are studied principles of rock mechanics (stress, strain and rheology).

- Learning outcomes and competences

Knowledge and understanding of the fundamental concepts of structural geology and its applications. Development of practical skills in the field of structural mapping and graphical techniques for representation and analysis of structural data.

- Teaching methods and learning activities

In the classes we use presentations like PowerPoint, which include graphs, diagrams, images, photographs and field photos. These materials are placed in one file early in the semester, so students can organize their files in advance. We also use Flash videos related to the processes of structures formation.

Observation of hand samples is also performed. A field trip is always done in the end of the semester.

In some classes exercises which involve theoretical concepts are presented.

- Structural geology

Is the study of the three-dimensional distribution of rock units with respect to their deformational history?

- What is structural geology?

Structural Geology Course for 3rd Year Students - Geology Department

Study of rock deformation; i.e. “the study of the architecture of the Earth’s crust”.

- **Structure Vs Tectonics**

- **Definition**

Structure comes from Latin word “Struere”, which means “to build”.

Structural geology is concerned with the reconstruction of the motions that have shaped the evolution of the Earth’s outer layer.

Tectonics (large scale structures) comes from Greek word “*Tektos*” which means “builder”.

- **Structural geology is the study of:**

“Structural Geology” is the study of the deformation of rocks and its effects.

The geometry of geologic structures (such as faults, folds, joints, intrusive bodies, etc.)

How these structures form (their relationship to applied stress)

Their significance to the geologic history of an area.

Their relationship to plate-tectonic motions.

- **Goals of structural geology**

This course aims at:

- Study of rock masses in 3-D
- How rocks are deformed?
- How structures are formed?
- The primary goal of structural geology is to use measurements of present-day rock geometries to uncover information about the history of deformation.

A common goal is to understand the structural evolution of a particular area with respect to regionally widespread patterns of rock deformation


▪ **Importance of structural geology**

The study of geologic structures has been of prime importance in economic geology, both petroleum geology and mining geology. Folded and faulted rock strata commonly form traps that accumulate and concentrate fluids such as petroleum and natural gas. Similarly, faulted and structurally complex areas are notable as permeable zones for hydrothermal fluids, resulting in concentrated areas of base and precious metal ore deposits. Veins of minerals containing various metals commonly occupy faults and fractures in structurally complex areas. These structurally fractured and faulted zones often occur in association with intrusive igneous rocks. They often also occur around geologic reef complexes and collapse features such as ancient sinkholes. Deposits of gold, silver, copper, lead, zinc, and other metals, are commonly located in structurally complex areas.

Structural geology is a critical part of engineering geology, which is concerned with the physical and mechanical properties of natural rocks. Structural fabrics and defects such as faults, folds, foliations and joints are internal weaknesses of rocks which may affect the stability of human engineered structures such as dams, road cuts, open pit mines and underground mines or road tunnels.

Environmental geologists and hydrogeologists need to apply the tenets of structural geology to understand how geologic sites impact (or are impacted by) groundwater flow and penetration. For instance, a hydrogeologist may need to determine if seepage of toxic substances from waste dumps is occurring in a residential area or if salty water is seeping into an aquifer.

- **Geological Structures**




- **Applications of Structural Geology**

Mineral resources

Crystallization in fluid-rich environment

Hot, metal-rich fluid migration as **vein deposits** or **disseminated deposits** (remobilized by faults)


(Quartz Veins)




(modified from slide by Ramon Arrowsmith)

Oil Traps

Oil Traps



Types of oil traps
(Keller, 2002)



▪ **STRESS Vs Strain**

STRESS


Study of the action and effect of forces on bodies.

STRESS

Stress [S] Force per unit area or the intensity of Force

$$S = F/A$$

Stress = (Force / Area)



▪ **Strain**

Strain = A change in shape or volume in response to a stress.

Stress is the cause, strain is the effect!



Stress Axes

σ_1 (Maximum Principal Stress Axis)

σ_2 (Intermediate Principal Stress Axis)

σ_3 (Minimum Principal Stress Axis)

(Hobbs et al., 1976) ($\sigma_1 \geq \sigma_2 \geq \sigma_3 \neq 0$)

The behavior of rock during stress

Depends on:

- 1) The type (tension, compression, shear) and amount of stress.
- 2) The temperature.
- 3) The pressure.
- 4) The rate at which the stress is applied

(Think about Silly Putty).


Rock squeezing experiments reveal 3 types of rock behavior:

- 1) Elastic behavior
- 2) Brittle behavior
- 3) Ductile/plastic behavior




Elastic Behavior – A lightly-deformed material will recover its original shape after the stress is removed. Elastic behavior occurs at very low stresses.

Every material has an **elastic limit**, the maximum amount of stress a material can feel and still recover to its original shape




The behavior of rock after reaching the elastic limit depends on T and P.

Brittle Behavior – Near the Earth's surface (low T and P), rocks exhibit brittle behavior...rocks will fracture (faults).




Ductile Behavior

At depth in the Earth (high T and P), rocks exhibit plastic behavior.... rocks will bend and flow (folds).




21

Strength of rock



Evidence of Crustal Deformation

- Folding of strata
- Faulting of strata
- Tilting of strata
- Joints and fractures




Evidence of Crustal Deformation

- Folding of strata
- Faulting of strata
- Tilting of strata
- Joints and fractures




How do you read this map?



- How do you read this map?

Structural Geology Course for 3rd Year Students - Geology Department



- How do you explain this structure?



- How do you read this map?

Chapter- II

Tectonics

▪ Introduction

- In 1785, James Hutton introduces; **(the present is the key to the past).**

A group of scientists started to recognize themselves as geologists. Their main aims were:

- To make geological maps.
- Reported the formation of rocks.
- The origins of specific structures and mountain ranges.

Later, ideas about the origin of mountains have evolved gradually.

Some questions we will answer

- How is the earth always changing?
- What forces inside the earth create and change landforms on the surface?
- What is continental drift and sea floor spreading?
- What is the theory of plate tectonics and how does it work?
- What two theories help make up the theory of plate tectonics?


What happens when the plates crash together, pull apart, and slide against each other?

▪ The Earth's crust


- The Earth is made of many different and distinct layers. The deeper layers are composed of heavier materials; they are hotter, denser and under much greater pressure than the outer layers.

Structural Geology Course for 3rd Year Students - Geology Department


- Natural forces interact with and affect the earth's crust, creating the landforms, or natural features, found on the surface of the earth.



- Before we start to look at the forces that contribute to landforms, let's look at the different layers of the earth that play a vital role in the formation of our continents, mountains, volcanoes, etc.



A slice through the globe to show Earth's different layers



crust - the rigid, rocky outer surface of the Earth, composed mostly of basalt and granite. The crust is thinner under the oceans.

mantle - a rocky layer located under the crust - it is composed of silicon, oxygen, magnesium, iron, aluminum, and calcium. Convection (heat) currents carry heat from the hot inner mantle to the cooler outer mantle.

outer core - the molten iron-nickel layer that surrounds the inner core.

inner core - the solid iron-nickel center of the Earth that is very hot and under great pressure.


393

Continental Drift Theory and Seafloor Spreading


▪ Continental Drift Theory

- When the tectonic plates under the continents and oceans move, they carry the continents and oceans with them.
- In the early 1900s a German explorer Alfred Wegner; proposed the **continental drift theory**. He proposed that there was once a single “supercontinent” called Pangaea.

Structural Geology Course for 3rd Year Students - Geology Department




Wegner's theory was that about 180 million years ago, Pangaea began to break up into separate continents. To back this theory up, he preserved remains and evidence from ancient animals and plants from South America, Africa, India, and Australia that were almost identical.




▪ Seafloor Spreading

The other theory supporting plate tectonics emerged from the study of the ocean floor.

Scientists were surprised to find that rocks taken from the ocean floor were much younger than those found on the continents. The youngest rocks were those nearest the underwater ridge system which is a series of mountains that extend around the world, stretching more than 64 thousand kilometers (40 thousand miles).




Seafloor Spreading



The theory of seafloor spreading suggests that molten rock (think of a melted chocolate bar that has been left in your pocket for too long)... This hot substance (lava) from the mantle rises under the underwater ridge and breaks through a split at the top of the ridge (the crust... Remember, the plate). The split is called a rift valley. The rock then spreads out in both directions from the ridge as if it were on two huge conveyor belts. As the seafloor moves away from the ridge, it carries older rocks away. Seafloor spreading, along with the continental drift theory, became part of the theory of plate tectonics.

Plate motions also can be looked at into the future, and we can have a stab at what the geography of the planet will be like. Perhaps in 250 million years time there will be a new supercontinent.



- What two theories help make up the theory of plate tectonics?
- What is continental drift and sea floor spreading?


▪ Plate Tectonics

- Most of these changes in the earth's surface takes place so slowly that they are not immediately noticeable to the human eye.
- The idea that the earth's landmasses have broken apart, rejoined, and moved to other parts of the globe forms part of the plate tectonic theory.


About 60 years ago, scientists exploring the seafloor found that it is full of tall mountains and deep trenches, a single seafloor mountain chain circles Earth and contains some of Earth's tallest mountains.

Along this mountain chain is a deep crack in the top layers of earth. Here the seafloor is pulling apart and the two parts are moving in opposite directions, carrying along the continents and oceans that rest on top of them. These pieces of Earth's top layer are called *tectonic plates*. They are moving very slowly, but constantly. (Most plates are moving about as fast as your fingernails are growing -- not very fast!) Currently Earth's surface

layers are divided into nine very large plates and several smaller ones.



According to the theory of plate tectonics, the earth's outer shell is not one solid piece of rock. Instead the earth's crust is broken into a number of moving plates. The plates vary in size and thickness.



The North American Plate stretches from the mid-Atlantic Ocean to the northern top of Japan. The Cocos Plate covers a small area in the Pacific Ocean just west of Central America.

These plates are not anchored in place but slide over a hot and bendable layer of the mantle.

In Conclusion:

Plate Tectonics

Plate Tectonic Theory:

Plate Tectonic Theory says that; the earth's crust and the upper mantle are divided into major lithospheric plates moving tangentially with respect to each other.

But this doesn't actually tell me how the mountains or volcanoes were formed or how earthquakes happen, does it?


YES!

- As mentioned earlier, those tectonic plates are always moving. They are always moving:
 - pulling away from each other
 - crashing head-on
 - or sliding past each other.

Depending on which way these plates are moving will decide what is happening on the earth you and I are standing on.

They're Pulling Apart!

- When plates pull away from one another they form a diverging plate boundary, or spreading zone.




Thingvellir, the spreading zone in Iceland between the North American (left side) and Eurasian (right side) tectonic plates. *January 2003.*

What happens when plates crash into each other depends on the types of plates involved.


Because continental crust is lighter than oceanic crust, continental plates "float" higher.

Therefore, when an oceanic plate meets a continental plate, it slides under the lighter plate and down into the mantle. The slab of oceanic rock melts when the edges get to a depth which is hot enough. (A temperature hot enough to melt silicon about a thousand degrees!) This process is called subduction. Molten material produced in a subduction zone can rise to the earth's surface and cause volcanic building, mountains, and islands.



What are Plates?


- The *Earth's crust and upper mantle (Lithosphere)* are broken into sections called plates




Plates move around on top of the mantle like rafts

A section of the lithosphere that slowly moves over the asthenosphere, carrying pieces of continental and oceanic crust.

What makes the plates move?




Convection Currents in the mantle move the plates as the core heats the slowly-flowing asthenosphere (the elastic/plastic-like part of the mantle).



▪ **Plate Boundaries**

The edges of Earth's plates meet at plate boundaries.

❖ Extended deep into the lithosphere



FAULT – Breaks in Earth's crust where rocks have slipped past each other.

THERE ARE THREE TYPES OF PLATE BOUNDARIES!


What are the three types of boundaries?

- **Divergent Boundaries**
- **Convergent Boundaries**
- **Transform Boundaries**


A different type of plate movement occurs along each type of boundary.

Divergent Boundaries


A plate boundary where two plates move away from each other.



RIFTING
causes
SEAFLOOR
SPREADING



Oceanic-Continental Collision



How is the rock *pulled* at Divergent Boundaries?

Rock gets THIN in the middle as it is pulled apart.




This STRESS is called **Tension**


What happens next at Divergent Boundaries?

- A geologic feature or event...

May form RIFT VALLEYS on continents





SEA-FLOOR SPREADING in the ocean




Features of Divergent Boundaries

- Mid-ocean ridges
- rift valleys
- fissure volcanoes




Convergent Boundaries




There are 3 types of Convergent Boundaries...

Type 1




What else happens at Convergent Boundaries?


VOLCANOES
occur at
subduction
zones



Type 2

- **Ocean plate** colliding with another **ocean plate**
- The less dense plate slides under the more dense plate creating a **subduction zone** called a **TRENCH**






Aleutian Islands, Alaska


Type 3

- A continental plate colliding with another continental plate
- Have Collision Zones:
 - A place where folded and thrust faulted mountains form.




- May form **Mountain Ranges**.

These are Folded Mountains, like the Himalayas or the Rockies.




Continental-continental Collision



(c)

Transform Boundaries

A plate boundary where two plates move past each other in opposite direction.



How is the rock broken at Transform Boundaries?

- Rock is pushed in two **opposite directions** (or sideways, but no rock is lost)
- This stress is called **SHEARING**



Transform Fault

What happens next at Transform Boundaries?


- May cause **Earthquakes** when the rock snaps from the pressure.
- A famous fault @ a Transform Boundary is the *San Andreas Fault in California.*




▪ **HOTSPOTS**

Map of part of the Pacific basin showing the volcanic trail of the Hawaiian hotspot-- 6,000-km-long Hawaiian Ridge-Emperor Seamounts chain.


A sharp bend in the chain indicates that the motion of the Pacific Plate abruptly changed about 43 million years ago, as it took a more westerly turn from its earlier northerly direction. Why the Pacific Plate changed direction is not known, but the change may be related in some way to the collision of India into the Asian continent, which began about the same time.




OTHER HOTSPOTS




■ Plate Motion Summary



▪ **Tectonic setting of the Red Sea**





Tectonic setting of the Red Sea. Modified from Bosworth et al. (2005).
 Red arrows are GPS-derived velocities from ArRajehi et al. (2010).




Regional setting of the Red Sea, Gulf of Suez and Gulf of Aqaba. Instrumentally recorded earthquakes demarcate the plate boundaries of the region (M 4 since 1960-01-01; ANSS 2016). Yellow arrows show the slip rates of Arabia and Sinai relative to Africa (ArRajehi et al. 2010). The small arrow on Sinai equates to 0.15 cm/yr (Mahmoud et al. 2005). Locations of Figs. 2 and 14 are shown by boxes. Volcanic rocks of Harrat Al Birk are shown in red. The continental Danakil Block is shown in green.

- **East African Rift System**

East African Rift System, also called **Afro-Arabian Rift Valley**, one of the most extensive rifts on Earth's surface, extending from Jordan in southwestern Asia southward through eastern Africa to Mozambique. The system is some 4,000 miles (6,400 km) long and averages 30–40 miles (48–64 km) wide.



East African Rift System



Chapter -III

Primary Structures

Primary structures

A primary structure is defined by Wilkerson (2019) as, "any structure that develops prior to or during the formation of the rock." Primary structures are non-tectonic, meaning they form during sedimentary deposition, or in the case of igneous rock, during crystallization.

We will explain primary structures by grouping them into five subcategories:

- Depositional Structures
- Deformational Structures
- Sedimentary Structures
- Igneous Structures
- Depositional Contacts

Primary structures are useful to geologists for many reasons. For one, they can be used to determine facing, or which side of the rock represents the upward direction during deposition. Primary structures can also be used to describe the strain the rock has undergone, as well as adding details regarding the geologic setting during deposition.

Sedimentary structures including:

- **Bedding** – Primary layering in a sedimentary rock, formed during deposition, manifested by changes in texture, color, and/or composition; may be emphasized in outcrop by the presence of parting
- **Overtured beds** – Beds that have been rotated past vertical in an Earth–surface frame of reference; as a consequence, facing is down
- **Parting** – The tendency of sedimentary layers to split or fracture along planes parallel to bedding; parting may be due

Structural Geology Course for 3rd Year Students - Geology Department

to weak bonds between beds of different composition, or may be due to a preference for bed-parallel orientation of clay
Bedding = layering = stratification




Bedding – Primary layering




Sedimentary structures

Stratigraphic facing (= Younging direction) – The direction to younger strata, or, in other words, the direction to the depositional top of beds.




- **The use of bedding in structural analysis**
 - The Law of Original Horizontality
 - Bedding provides a reference frame.
 - Bedding is labeled S_0 (pronounced ess-zero), where the S is an abbreviation for planar structures (surface)
 - Bedding provides information on depositional environment, younging direction, current direction.
 - Homoclinal.

- **Graded beds and younging direction**
 - During settling, the largest grains fall first, and the finest grains last
 - Bouma sequence in turbidites
 - Flysch



Cross beds and younging direction

- Sedimentation on a lee side
- Topset, forset, and bottomset beds
- Erosion of topset beds
- Clear stratigraphic facing and current indicator





Cross bedding

- **Convolute folds**



▪ **Disrupted Bedding**



▪ **Disrupted Bedding**

- Load casts
- Sand volcanoes
- Clastic dikes

Studies of disrupted bedding, sedimentary dikes, and sand volcanoes provide an important basis for determining the recurrence interval of large earthquakes

Load casts


Extend downward from a sand layer into an underlying mud or very fine sand layer



Load casts



Load casts



Clastic dikes



▪ **Conformable and unconformable contacts**


Disconformity beds above and below the unconformity are parallel to one another, but there is an age difference between the two sequences.

Angular unconformity strata below the unconformity have a different attitude than strata.

Nonconformity strata were deposited on a basement of older crystalline rocks.

Structural Geology Course for 3rd Year Students - Geology Department

Buttress unconformity beds of the younger sequence were deposited in a region of significant predepositional topography




Angular unconformity




Nonconformity




How to identify unconformities?




(a)



(b)



(c)



(d)

Structural Geology Course for 3rd Year Students - Geology Department

- **Compaction and Diagenetic Structures**
Compaction results in a decrease in porosity (>50% in shale and >20% in sand) that results in an increase in the density of the sediment
- Compaction of mud leads to development of a preferred orientation of clay - shale
- Deeper compaction can cause pressure solution

- **Pen contemporaneous Structures**
 - Deposition on a preexisting slope or tilting prior to full lithification in a tectonically active region can pull the layers down the slope
 - Fluid pressure in the layers keeps the layers apart
 - Debris flows
 - The deformed interval is *intraformational*, meaning that it is bounded both above and below by relatively unreformed strata




Chapter- IV

Faults

- Introduction

- **FAULTING**

Deformation Styles



Brittle Vs Ductile

Brittle

- Grain size reduction by brittle fracture
- Randomly oriented fabrics
- Fluid flow + cementation/silicification
- Frictional melting (pseudotachyllite)

Ductile

- Grain size reduction by crystal plastic strain
- Systematically oriented fabrics
- Development of foliation
- Development of stretching lineation


Make sure that you understand the big picture

The behavior of rock during stress

Depends on:

- 1) **The type (tension, compression, shear) and amount of stress.**
- 2) **The temperature.**
- 3) **The pressure.**
- 4) **The rate at which the stress is applied**
(Think about Silly Putty).

Make sure that you understand the big picture!



- **BRITTLE deformation**

Brittle deformation of an object refers to deformation promoting loss of continuity of or within the object. The discontinuities eventually created during brittle deformation are called fractures. In contrast to ductile deformation, which implies internal continuity of the object after deformation, brittle deformation occurs mostly in the upper structural levels of the crust.

We will use the word *fracture* to characterize a brittle structure without any other precision, i.e. whenever details on the relative


Structural Geology Course for 3rd Year Students - Geology Department

displacements of the fracture walls cannot be assessed. As such 'fracture' is used as a general term for all kinds of brittle structures. Fractures are in turn grouped in *mode I, II and III fractures*, depending on the relative motions of their fracture walls.

Brittle deformation; the permanent change that occurs in a solid material due to the growth of fractures and/ or due to sliding on fractures. Brittle deformation only occurs when stresses exceed a critical value, and thus only after a rock has already undergone some elastic and/or plastic behavior.

Brittle fault zone A band of finite width in which slip is distributed among many smaller discrete brittle faults, and/or in which the fault surface is bordered by pervasively fractured rock.

brittle fault offsets



- **DUCTILE deformation**


- Ductile deformation mechanisms

Ductile deformation indicates shape change of a material through bending or flowing during which chemical bonds may become broken but subsequently reformed into new bonds. It requires stress surpassing the elastic threshold and a deformation rate slow enough to accommodate further strain without breaking of the material

Structural Geology Course for 3rd Year Students - Geology Department

(i.e., ductility or plasticity). Fold, foliation, and lineation are the typical features of a rock that experienced ductile deformation, though the last two can sometimes also be formed by brittle deformation. Several mechanisms are responsible for ductile deformation, including diffusion creep, dislocation creep, mechanical twinning/kinking, grain boundary sliding, and rigid body rotation.

Ductile offsets within shear zone



another example of a small ductile shear zone




▪ **FAULTS**

A fault is a planer fracture across which the rock has been displaced in a direction which is generally parallel to the fracture plane.

Faults are ruptures along which the opposite walls have moved past each other. The essential features is differential movement parallel to the surface of the fracture.


• **Fault geometry**



Strike-slip and dip-slip components

Oblique-slip faults

- Strike-slip and dip-slip components
- Most faults are oblique-slip, but are often dominantly strike-slip or dip-slip




Geometry of displacement

- The faults with displacement parallel to the strike of the fault plane are termed **Strike-slip faults** . These faults may also be called **Wrench, Tear or Trans current faults**.

The faults with displacement parallel to the dip of the fault plane are termed **dip-slip faults** .

- Faults with oblique-slip displacement are regarded as having strike-slip and dip-slip components are termed **oblique faults** .



The Main Types of Faults

Normal Fault

Normal Fault; Qena – Safaga Road



Thrust Fault

Thrust Fault; Qena-Safaga Road




Strike-Slip Fault


Strike-slip Faults; Qena- Safaga Road



Fault - Fault zone - Shear zone

Fault - Fault zone - Shear zone







What are the three kinds of faults?

- The three main kinds of faults are *strike-slip faults*, *normal faults*, and *reverse faults*.
- A *fault plane* is the location where two fault blocks meet.

For any fault except a perfectly vertical fault, the block above the fault plane is called the *hanging wall*. The block below the plane is the *footwall*.




▪ Strike-slip faults

- In a strike-slip fault, the fault blocks move past each other horizontally.
- Strike-slip faults form when rock is under **shear stress**, or stress that pushes rocks in parallel but opposite directions.


Strike-slip faults are common along transform boundaries, where tectonic plates move past each other

• What Governs Strike-Slip Fault Movement


Geologists understand the mechanics of strike-slip faulting at the plate scale but up until now, the specifics of faulting on a very small scale haven't been as thoroughly studied. Often times, we see faults as they currently exist but don't get to look at the development of that fault from incipient stages.



- **Dextral and sinistral strike- slip faults**



- **Right – lateral [Dextral] Strike- slip Fault**



- **example right – lateral [dextral] strike- slip Fault**



- **SAN ANDREAS**

- **WHAT TYPE OF FAULT IS THE SAN ANDREAS?**

A San Andreas earthquake would be classified as occurring on a strike-slip fault. Strike-slip faults are found along boundaries of tectonic plates sliding past each other.

A strike-slip fault is a vertical fracture in the earth's crust that creates horizontal motion, along the line of the fault. The walls of rock move to the left relative to one another, or to the right relative to one another. These faults are formed by horizontal compression.

Motion along a strike-slip fault is horizontal.

Shear stress is acting on the opposing earthen blocks.

The fault surface is near vertical.

Stress on the San Andreas fault causes right-lateral motion



Example

Right-lateral
Strike-slip Fault

San Andreas Fault


The **San Andreas Fault** is a continental [transform fault](#) that extends roughly 1,200 kilometers (750 mi) through [California](#). It forms the [tectonic](#) boundary between the [Pacific Plate](#) and the [North American Plate](#), and its motion is [right-lateral strike-slip](#) (horizontal). The fault divides into three segments, each with different characteristics and a different degree of earthquake risk. The slip rate along the fault ranges from 20 to 35 mm (0.79 to 1.38 in)/yr. It was formed by a transform boundary.




extends roughly 1,200 kilometers (750 mi) through [California](#). It forms the [tectonic](#) boundary between the [Pacific Plate](#) and the [North American Plate](#), and its motion is [right-lateral strike-slip](#); and extends to depths of at least 10 miles within the Earth

Arrows show relative motion of the North American Plate (southeastward) and the Pacific Plate (northwestward)

- **Dead Sea Fault**




- **Left – lateral [sinistral] Strike- slip Fault**



- **Najd fault system**

The Najd Fault System is a complex set of left-lateral strike-slip faults and ductile shear zones that strike NW-SE across the Precambrian of Arabia and Egypt. This system was developed during the interval 540–620 Ma. It is up to 400 km wide with an exposed length of 1100 km; inferred buried extensions of the Najd give it a total length of 2000 km. It is the best exposed and may be the largest pre-Mesozoic zone of transcurrent faulting on earth. Previous models for the Najd Fault System suggest it formed as a result of a major Late Precambrian continent-continent collision. This model is not preferred here because (1) the lack of evidence for a pre-Late Precambrian continent to the east of the Najd Fault System; (2) the difference between the orientation of the Najd Fault System and that predicted by slip-line theory; (3) the younger age of Najd movements compared with that of collisional sutures in the Arabian Shield; and (4) lack of evidence for wide-spread crustal uplift that would be expected to accompany collision. A new model for the origin of the Najd Fault System accounts for each of these objections: The Najd Fault System formed in response to a broad zone of NW-SE directed crustal extension that accompanied juvenile continental crustal formation in northernmost Afro-Arabia. This model also accounts for the following observations: (1) Strands of the Najd parallel the direction of extension in the North Eastern Desert of Egypt and Sinai; (2) the timing of the principal rifting movements (ca. 575–600 Ma) overlap with those of the Najd (ca. 560–620 Ma); (3) in spite of observation (2), the Najd Fault System is not recognized in northernmost Afro-Arabia; instead the Najd deformation becomes increasingly ductile and these zones are more commonly intruded by sheared and foliated granites as the principal zone of extension is approached. The Najd Fault System thus represents a set of continental transforms developed in response to a major episode of Late Precambrian extensional continental crust formation in northernmost Afro-Arabia





Najd fault system and sutures of the northern part of the Arabian-Nubian Shield. Black arrows show principal stress directions for the Najd fault system as a whole. Small white arrows show strain in individual gneiss complexes. Note that this is extensional in the Eastern desert, but transpressive in the Feiran–Solaf metamorphic complex. (Compiled from various sources, including: Abdelsalam & Stern, 1996; Fritz et al., 1996; Johnson, 1998).

- **Pull-apart basins**

These basins are called pull-apart basins because the crust is literally pulled apart in the section between the two strike-slip faults.

Examples. Two famous localities for continental pull-apart basins are the **Dead Sea and Salton Sea**. Pull-apart basins are amenable to research because sediments deposited in the basin provide a timeline of activity along the fault.

The **Salton Sea** is a shallow, landlocked, highly saline body of water in Riverside and Imperial counties at the southern end of the U.S. state of California.



Pull-apart basins represent an important exploration target for oil and gas, [porphyry copper](#) mineralization, and [geothermal](#) fields. The Matzen fault system in the [Matzen oil field](#) has been recast as extensional [grabens](#) produced by pull-apart basins of the [Vienna Basin](#). The [Dead Sea](#) has been studied extensively and thinning of the crust in pull-aparts may generate differential loading and


instigate [salt diapirs](#) to rise,^[9] a frequent [trap](#) for hydrocarbons. Likewise intense deformation and rapid subsidence and deposition in pull-aparts creates numerous structural and stratigraphic traps, enhancing their viability as [hydrocarbon reservoirs](#).^[10]

The shallow extensional regime of pull-apart basins also facilitates the emplacement of [felsic intrusive rocks](#) with high [copper](#) mineralisation. It is believed to be the main structural control on the giant [Escondida](#) deposit in [Chile](#).^[11] [Geothermal](#) fields are located in pull-aparts for the same reason due to the high heat flow associated with rising magmas

• Reverse Faults

- In a reverse fault, the hanging wall moves up relative to the footwall.
- Reverse faults form when rocks undergo **compression**, which is stress that squeezes or pushes rock together.


Reverse faults are common along convergent boundaries, where two plates collide.




- **Thrust faults**

A **thrust fault** is a break in the Earth's crust, across which older rocks are pushed above younger rocks.

A thrust fault is a reverse fault with a dip of 45° or less, a very low angle.




- **Reverse Faults and thrust faults**




- **Overthrust fault**

Thrust fault is a type of reverse fault that has a dip of 45 degrees or less. If the angle of the fault plane is lower (often less than 15 degrees from the horizontal) and the displacement of the overlying block is large (often in the kilometer range) the fault is called an over-thrust or over-thrust fault.



- **Nappe Structure**

Nappes form when a mass of rock is forced (or "thrust") over another rock mass, typically on a low angle fault plane. The resulting structure may include large-scale recumbent folds, shearing along the fault plane, imbricate thrust stacks, fensters and klippe.




- **Window (Fenster)**

- Thrust faults are often thin sheets, and erosion may open holes in them
- A hole through a thrust sheet is called **a fenster**, or window
- **Fenster**: An eroded area of a thrust sheet that displays the rocks beneath the thrust sheet
- Triangular teeth point outward fenster are used on a map

Klippe

- If erosion leaves an isolated remnant of thrust sheet, surrounded by exposed footwall, the remnant is called a **klippe** (German for cliff)
- **Klippe** are indicated on a map by inward pointing teeth.

- **Bedding and Fault Plane Orientation**



Fault segments may parallel bedding in either the footwall of the hanging wall, but cut across bedding in the opposite block.

- **Fault-bend folds**

Thrust faults, particularly those involved in [thin-skinned](#) style of deformation, have a so-called *ramp-flat* geometry. Thrusts mostly propagate along zones of weakness within a sedimentary sequence, such as [mudstones](#) or [halite](#) layers, these parts of the thrust are called [decollements](#). The part of the thrust linking the two flats is known as a *ramp* and typically

forms at an angle of about 15°–30° to the bedding. Continued displacement on a thrust over a ramp produces a characteristic fold geometry known as a *ramp anticline* or, more generally, as a *fault-bend fold*




Diagram of the evolution of a fault-bend fold or 'ramp anticline' above a thrust ramp, the ramp links [decollements](#) at the top of the green and yellow layers


- **Thrust duplex**

Duplexes occur where there are two decollement levels close to each other within a sedimentary sequence, such as the top and base of a relatively strong [sandstone](#) layer bounded by two relatively weak mudstone layers. When a thrust that has propagated along the lower detachment, known as the *floor thrust*, cuts up to the upper detachment, known as the *roof thrust*, it forms a ramp within the stronger layer. With continued displacement on the thrust, higher stresses are developed in the footwall of the ramp due to the bend on the fault. This may cause renewed propagation along the floor thrust until it again cuts up to join the roof thrust. Further displacement then takes place via the newly created ramp. This

process may repeat many times, forming a series of fault bounded thrust slices known as *imbricates* or [horses](#), each with the geometry of a fault-bend fold of small displacement. The final result is typically a lozenge shaped duplex.

Most duplexes have only small displacements on the bounding faults between the horses and these dip away from the foreland. Occasionally the displacement on the individual horses is greater, such that each horse lies more or less vertically above the other, this is known as an *antiformal stack* or **imbricate stack**. If the individual displacements are greater still, then the horses have a foreland dip.

Duplexing is a very efficient mechanism of accommodating shortening of the crust by thickening the section rather than by folding and deformation.



Development of thrust duplex by progressive failure of ramp footwall


- **Blind thrust faults**

If the fault plane terminates before it reaches the Earth's surface, it is referred to as a *blind thrust* fault. Because of the lack of surface evidence, blind thrust faults are difficult to detect until they rupture.


The destructive [1994 earthquake in Northridge, Los Angeles, California](#), was caused by a previously undiscovered blind thrust fault.

Because of their low [dip](#), thrusts are also difficult to appreciate in mapping, where lithological offsets are generally subtle and stratigraphic repetition is difficult to detect, especially in [peneplain](#) areas.


- **Examples of thrust belts**



Zagros thrust belt



- **Zagros thrust belt**





- **Zagros thrust belt**

The collision of the Iranian microcontinent with the Afro-Arabian continent resulted in the deformation of the Zagros [orogenic belt](#). The foreland of this belt in the [Arabian Gulf](#) and Arabian platform has been investigated for its petroleum and gas resource

potentials, but the Zagros hinterland is poorly investigated and our knowledge about its deformation is much less than other parts of this orogen. Therefore, this work presents a new geological map, stratigraphic column and two detailed geological cross sections


▪ Normal faults

- In a normal fault, the hanging wall moves down relative to the footwall, in a way you would *normally* expect as a result of gravity.
- Normal faults form when rock is under **tension**, which is stress that stretches or pulls rock apart.
- Normal faults are common along divergent boundaries.




Normal fault


• Horst Vs Graben




Horst and graben structure, typical rift related structure (direction of extension shown by red arrows)



Schematic evolution of low-angle normal faults. (A) Under normal crustal conditions, faults rotate via domino-style block rotation (after Fletcher and Spelz, 2009). In this model, multiple faults form at high angles over a distributed region and simultaneously begin to rotate to shallower angles (e.g., Chamberlin, 1983; McClay and Ellis, 1987; Whitney et al., 2013). Faults are eventually crosscut by a younger generation of faults after they have rotated $\sim 20^\circ$ - 45° from their initial orientations (Nur et al., 1986). (B) Under hot crustal conditions, faults rotate via a rolling-hinge mechanism (Buck, 1988; Wernicke and Axen, 1988; Lavier et al., 1999). In this model, slip is concentrated along a single large-offset fault system. As extension progresses, isostatic uplift rotates the initial fault to shallower angles, and new faults emerge basinward as older faults are continually rotated.





Structural setting and tectonic evolution of the Gulf of Suez





- **Listric Fault**

Listric Fault: In the field of geology, a listric fault refers to **a fault with a curved fault plane**. Most listric faults comprise a steeply dipping section near the surface. But the fault plane becomes increasingly flat with depth. Listric faults can be normal faults or reverse faults





Listric normal fault with a family of rotating "domino" blocks bounded by synthetic, planar faults (from Wernicke & Burchfiel 1982).



• **Stress distribution during faulting**


$\sigma_1 > \sigma_2 > \sigma_3$
 σ_1 vertical; extension
 σ_1 diagonal; strike-slip
 σ_1 horizontal; Thrust



- **FAULT CRITERIA**

- *Displacement of recognizable marker such as fossils, color, composition, texture, .etc.).*
- *Repetition or omission of stratigraphic units asymmetrical repetition*
- *Occurrence of fault rocks (mylonite or cataclastic or both)*

- *Presence of S or C structures or both, rotated porphyry clasts and other evidence of shear zone.*
 - *Abundant veins, silicification or other mineralization along fracture may indicate faulting.*
 - *Drag Units appear to be pulled into a fault during movement (usually within the drag fold and the result is thrust fault)*
 - *Reverse drag occurs along listric normal faults.*
 - *Slickensides and slickenlines along a fault surface*
 - *Topographic characteristics such as drainages that are controlled by faults and fault scarps.*
-
- **Recognition of fault in the field**
 - Discontinuity of structures.
 - Repetition or omission of strata.
 - Presence of features characteristic of fault-planes (slickenside, gouge, fault breccia)
 - Solidification and mineralization
 - *Displacement of recognizable marker*




Fault

Criteria for Faulting

2. Repetition or omission of strata

Repetition or omission of strata may indicate faulting. The following figure represents a geological map of a region of folded and faulted sedimentary rocks. A syncline lies near the centre of the map, as is shown by the dips, and the formations are progressively younger from a to e. In certain places, however, one or more formations are missing, as, for example, along the line FF, where formation b is absent, and along the line F'F', where c and d are missing. The lines FF and F'F' must be the traces of faults.



The omission of strata, however, may be due to an unconformity.




Criteria for Faulting

Fault

3. Features characteristics of fault planes

c. Drag : Drag is in some cases an aid in divulging the relative motion along the fault. Because of friction the beds in the up throw block are dragged up, whereas the beds in the down throw are drag down.




Department of Geology, University of Dhaka Elementary Structural Geology Slide 10 of 30  ©SHA


Criteria for Faulting

Fault

3. Features characteristics of fault planes

a. Slikensides:




Department of Geology, University of Dhaka Elementary Structural Geology Slide 8 of 30  ©SHA

Tension Gashes



Tension Gashes



Left-lateral motion

Features on the fault surface

- **Grooves** (parallel to the movement direction)
- **Growth of fibrous minerals** (parallel to the movement direction)
- **Slickensides** are the polished fault surfaces.
- **Small steps.**

All are considered a kind of lineation. They indicate the movement relative trend NW, NE ... etc.

Small steps may also be used to determine the movement direction and direction of movement of the opposing wall. Slicklines usually record only the last moment event on the fault.




Fault Rocks

SHEAR ZONE

Shear zones are produced by both homogeneous and inhomogeneous simple shear, or oblique motion and are thought of as zones of ductile shear.

Shear zones are classified by Ramsay (1980) as:

- 1) Brittle
- 2) Brittle-ductile
- 3) Ductile



• **Characteristics of Shear Zones**


Shear zones on all scales are zones of weakness.

- *Associate with the formation of mylonite.*
- *Presence of sheath folds.*
- *Shear zones may act both as closed and open geochemical systems with respect to fluids and elements.*
- *Shear zones generally have parallel sides.*

Displacement profiles along any cross section through shear zone should be identical.

INDICATORS OF SHEAR SENSE OF MOVEMENT


1. *Rotated porphyroblasts and porphyroclasts.*
2. *Pressure shadows*
3. *Fractured grains.*
4. *Boudins*
5. *Presence of C- and S- surfaces (parallel alignment of platy mineral)*
6. *Riedel shears.*



Examples of **ductile shear zones** in Piantonetto Valley. a Extremely localized, 'fault-like' shear zone. Pen (13.5 cm long) for scale. b Centimetre-thick approximately tabular shear zone with sharp boundary to the host metagranite. Coin (2 cm in diameter) .



Features from the Field: Shear Zones and Mylonites




Interpretation of the Neves shear zone, shown above. The foliation (yellow) forms in the metagranite close to the shear zone and it is deflected and parallelized to the mylonitic foliation (center). In the shear zone, the metagranite has been completely transformed to a mylonite. Photo © Samuele Papeschi © Samuele Papeschi

Why are shear zones important?


Understanding shear zones and interpreting their deformation history represent major goals for structural geologists. Fossil shear zones, now exhumed to the surface, hold valuable clues to how mountains formed and plates collided. They are precious 'hard drives' of the deformation mechanisms happening at depths and temperatures that are otherwise completely inaccessible. Studying those structures helps us understand how rocks deform and how strain is distributed through the lithosphere, giving us important clues about how plate tectonics works.

Structural Geology Course for 3rd Year Students - Geology Department

Shear zones also affect our daily lives! Even if shear zones are (generally, not always) aseismic (i.e. do not produce earthquakes), they transport mechanical energy loading shallower faults that have the potential to release energy in the form of destructive earthquakes. Shear zones are also major discontinuities that may act as preferred fluid pathways through the crust, becoming potential traps for strategic metal, rare earths, and other mineralizations.




- **S-C fabric**



S-C fabric is a metamorphic fabric formed by the intersection of shear surfaces within rocks affected by dynamic metamorphism

S-C fabric



S-C fabric is a metamorphic fabric formed by the intersection of shear surfaces within rocks affected by dynamic metamorphism. The foliation that develops in a shear zone is usually thought to trace the XY-plane of the strain ellipsoid. The sense of rotation of the foliation from the margin into the shear zone is generally a safe kinematic indicator. As strain accumulates, a set of slip surfaces or shear bands commonly forms parallel to the walls of the shear zone. These shear bands are called **C** (French *cisaillement* for shear, which relates to the movement of scissors) and the foliation is named **S** (for schistosity or *schistosité*).






FIGURE 12.11 Characteristic geometry of (a) C-S and (b) C-C' structures in a dextral shear zone. The C-surface is parallel to the shear zone boundary and is a surface of shear accumulation (i.e., not parallel to a plane of principal finite strain). The S-foliation is oblique to the shear-zone boundary and may approximate the XY-plane of the finite strain ellipsoid. The C'-foliation in (b) displaces an earlier foliation (C or composite C/S).



S-C fabric in a mylonite from Sardinia (Italy).
XPL image, 10x (Field of view = 2mm)

- **Fault Breccia**



Indurated Breccia Photomicrograph

- Note the angular fragments (fr) of quartz sandstone in a matrix of fine-grained iron oxide cement (ic)
- Field of View 4 x 2.7 mm, Cross Polarized Light

- Photomicrograph of fault breccia in the Antietam Formation, Blue Ridge province
- Breccias form when rocks are extensively fractured in fault zones and are cemented together when minerals precipitate in the cracks and fractures

• Mylonites

Mylonite is a fine-grained, compact [metamorphic rock](#) produced by [dynamic recrystallization](#) of the constituent [minerals](#) resulting in a reduction of the grain size of the rock. Mylonites can have many different [mineralogical](#) compositions; it is a classification based on the textural appearance of the rock.

Mylonites are [ductilely](#) deformed rocks formed by the accumulation of large [shear strain](#), in ductile [fault](#) zones. There are many different views on the formation of mylonites, but it is generally agreed that crystal-plastic deformation must have occurred, and that fracturing and cataclastic flow are secondary processes in the formation of mylonites. Mechanical abrasion of grains by milling does not occur, although this was originally thought to be the process that formed mylonites, which were named from the [Greek](#) *μύλος* *mylos*, meaning mill. Mylonites form at depths of no less than 4 km

• Blastomylonites

- *Blastomylonites* are coarse grained, often sugary in appearance without distinct [tectonic banding](#).
- *Ultramylonites* usually have undergone extreme grainsize reduction. In structural geology, ultramylonite is a kind of mylonite defined by modal percentage of matrix grains more than 90%. Ultramylonite is often hard, dark, [cherty](#) to [flinty](#) in

appearance and sometimes resemble [pseudotachylite](#) and [obsidian](#). In reverse, ultramylonite-like rocks are sometimes "deformed pseudotachylite".


- *Mesomylonites* have undergone an appreciable amount of grainsize reduction, and are defined by their modal percentage of matrix grains being between 50 and 90%.^{[9][10]}
- *Protomylonites* are mylonites which have experienced limited grainsize reduction, and are defined by their modal percentage of matrix grains being less than 50%. Because mylonitisation is incomplete in these rocks, relict grains and textures are apparent, and some protomylonites can resemble foliated [cataclasite](#) or even some [schists](#).

Porphyroclasts – a relict crystal in a ductile shear zone that has survived the shearing and reduction in grain size from the photolith ([Twiss & Moores, 1992](#)).

Porphyroblasts – mineral crystals which grow to relatively large sizes and are related to metamorphism and deformation ([Twiss & Moores, 1992](#))

Porphyroblasts / Porphyroclasts


- **Porphyroblasts** = large grains that have grown in the rock mass during deformation and metamorphism
- **Porphyroclasts** = relict earlier large grains that have survived ductile deformation (e.g. feldspar)



Blast: grain size increase
Clast: grain size decrease

Pseudotachylite

Pseudotachylyte




• Newer pseudotachylyte injection vein cuts the older one.

- Silicate rocks are excellent insulators, and heat generated by friction does not escape
- Temperatures in excess of 1000°C are possible
- Tachylyte is a type of volcanic glass, and the prefix *pseudo* means false, so the name literally means false volcanic glass

Foliation (mica) fish-


Foliation (mica) fish

- A layer with oblique internal fabric becomes detached along S-fabric to form an asymmetric 'fish'.




- Fish "flash" - oriented sample reflected sunlight...

- Tails on porphyroclasts are asymmetric in the direction of ductile flow.



Fractured Grains

- Large grains of some minerals (e.g. feldspars, pyroxenes) continue to deform brittlely within a ductile matrix.
- The fractures indicate a sense of motion opposite that of actual shear sense




How would you describe the kinematics of this system?

Death of a Fault

- Faults can also split, to form an anastomosing array, which may merge and diverge several times along its length
- A fault splay may develop, with the fault splitting and dying out – these are called horsetails (B)

- A fault dies when its displacement becomes less and less, finally reaching zero near the tip, in a zone of plastic deformation (C).



Chapter- V

Folds


Folds – warping of strata produced by compressive deformation, range in scale from microscopic features to regional-scale domes and basins.




▪ **Introduction**

- **Why we Study Folds?**

- **Folds can trap oil and natural.**




A



Folds are suitable sites for mineralization

- Folds are suitable sites for mineralization.

- Folds provide information about kinematic and deformational history of rock masses.




- **Conditions of Folding**

In structural geology, a fold is a stack of originally planar surfaces, such as sedimentary strata, that are bent or curved during permanent deformation. Folds in rocks vary in size from microscopic crinkles to mountain-sized folds. They occur as single isolated folds or in periodic sets (known as fold trains).

Folds form under varied conditions of stress, pore pressure, and temperature gradient, as evidenced by their presence in soft sediments, the full spectrum of metamorphic rocks, and even as primary flow structures in some igneous rocks.

Folds result from the slow deformation of rocks. This happens deep underground where the rocks are under pressure and temperatures are higher. Folded rocks are common in mountain ranges like the Alps, Himalayas and the Scottish Highlands



Main types of folds



Anticline: fold that is convex in the direction of the youngest beds

Syncline: Fold that is convex in the direction of the oldest beds



*requires that you know **facing direction** (direction of youngest beds); know stratigraphy!

Antiform: convex up

Synform: convex down
*simply describes geometry





▪ **Geometry of a Fold (Parts of a Fold)**



Folding – Geometric Description:

- The hinge connects the two limbs of a fold.
- The hinge point is the point of maximum curvature and is located in the center of the hinge zone.
- The hinge line is the three-dimensional equivalent of the hinge point.
- If the hinge line appears as a straight line it is called a **fold axis**.
- The axial surface (or **axial plane** when approximately planar) connects the hinge line of two or more folded surfaces.
- The axial trace represents the intersection between the axial plane and the surface of observation.
- The inflection point (inflection line) is where there is a change in curvature of a fold limb.
- The interlimb angle is the angle enclosed by the two limbs of a fold.
- The enveloping surface is the surface tangent to individual hinges along a folded layer.





▪ Classifications of Folds

Folds are generally classified according to the attitude of their [axes](#) and their appearance in cross sections perpendicular to the trend of the fold. The [axial plane](#) of a fold is the plane or surface that divides the fold as symmetrically as possible. The axial plane may be vertical, horizontal, or inclined at any intermediate angle. An axis of a fold is the intersection of the axial plane with one of the strata of which the fold is composed. Although in the simpler types of folds the axis is horizontal or gently inclined, it may be steeply inclined or even vertical. The angle of inclination of the axis, as measured from the horizontal, is called the [plunge](#). The portions of the fold between [adjacent](#) axes form the flanks, limbs, or slopes of a fold.

An anticline is a fold that is convex upward, and a syncline is a fold that is concave upward. An anticlinorium is a large anticline on which minor folds are superimposed, and a synclinorium is a large syncline on which minor folds are superimposed. A symmetrical fold is one in which the axial plane is vertical.

Structural Geology Course for 3rd Year Students - Geology Department

An asymmetrical fold is one in which the axial plane is inclined. An overturned fold, or overfold, has the axial plane inclined to such an extent that the strata on one limb are overturned. A recumbent fold has an essentially horizontal axial plane. When the two limbs of a fold are essentially parallel to each other and thus approximately parallel to the axial plane, the fold is called isoclinal.


▪ Anticline and Syncline

Older rocks will be in the core of an anticline. (An easy way to remember this is "antique" = old = *anti* cline = old rock will be in center.)

Younger rocks will be in the core of a syncline. (An easy way to remember this is "sin" = young = *syn* cline = younger rock will be in the center.)

Rock layers will dip down perpendicular in direction of the youngest rock at the surface.

The steeper a layer of rock dips the more narrow the width of its exposure belt and a wider dip represent a gentle dip.




- Anticline



-

- Syncline





Homocline

Homoclinal fold : is a region of uniform dip in one direction.




Fan folds : in which the two limbs are overturned.




Chevron folds : have sharp crests and troughs.




Box folds : the fold has two hinges and flat crest.




- Domes and Basins



Symmetric fold




Asymmetric fold




▪ Facing and vergence

The direction of apparent movement of the upper, long limb with respect to the shorter limb of an asymmetric fold is called the vergence. In other word, vergence is simply the sense of asymmetry. The true vergence or facing is the younging direction along the axial plane in a direction perpendicular to the fold axis. Uniformly verging asymmetric folds are characteristic of thrust belts. Vergence is useful in working out the regional direction of transport and help to fix an observation location on large folds.



2- Based on interlimb angle


2- Based on interlimb angle



Fold shape in the profile plane is described by the interlimb angle



Fold Profiles



3- Based on curvature of hinge zone

3- Based on curvature of hinge zone



Rounded fold



Angular fold

▪ Mechanism of Folding

Three distinct mechanisms have been identified for the folding of rocks: **bending**, **buckling**, and **passive folding**. Bending of rocks occurs when the deforming force is applied across (at high angle to) rock layers.


Folding Mechanisms

Types of folding:

- **Buckling** – active folding. There is a viscosity contrast between the folding layer and its host rock. Forces are applied parallel to the layer
- **Bending** – forces are applied across the layer.
- **Passive folding** – layers are simply passive markers with no rheological influence

The diagram illustrates three folding mechanisms. **Buckling** shows a horizontal layer being compressed from both ends by red arrows pointing towards each other. **Bending** shows a horizontal layer being pushed down from both ends by red arrows pointing downwards. **Passive folding** shows a layer being sheared from both ends by red arrows pointing in opposite directions, causing the layer to fold.


Passive – amplification of natural imperfections in the layers of a rock - or are a consequence of differential flow in a volume of rock.




Active (flexural folding) – the layering has mechanical significance

Buckle folding


occurs when there is a competency contrast between layers, and they are subjected to layer-parallel shortening.








Neutral surface folding:

Folded layer contains only one plane with no internal deformation, the neutral surface. Active folding.




Facing is very important in fold naming





En échelon folds

In some non-cylindrically folded surfaces, doubly plunging and relatively short folds with steeply dipping axial planes are arranged spatially such that culminations and depressions in successive folds lie along lines that make an acute angle with the axial surfaces. Such folds are stepped and consistently overlapping; they define an en échelon array. Note, however, that this term describes the geometry of the folded surface and is independent of the relationship of the structure to the horizontal and vertical.



Chapter –VI

FOLIATION and LINEATION

FOLIATION and LINEATION



▪ Introduction; "General" definition:

Foliation: Penetrative (at outcrop and microscopic scale) and parallel **planar** fabric elements in a rock.

"Structural geologist's" definition:

Foliation is a planar fabric, secondary and due to mineral recrystallization and/or plastic behavior during deformation at elevated temperatures

Gneissic structure:
Compositional banding
produced during
deformation.



Schistosity:
coarser grained
fabric- also a
foliation



Mylonitic foliation: Forms due to grain-size reduction by a mix of brittle and plastic deformation in shear zones





Figure 8.55 (A) The microscopic texture of mylonite. The white represents fragmented grains (porphyroclasts) and breccia streaks. The black represents plastically deformed quartz. (B) The microscopic texture of ultramylonite. [From Higgins (1971), Courtesy of United States Geological Survey.]

brittle deformation of feldspar porphyroclasts
plastic deformation of quartz "ribbons" and mica

Coarse-grained mylonitic augen gneiss. The large porphyroclasts are called augen ("eyes")



▪ Foliation Mechanism





FIGURE 5.1 Diagrammatic representation of various types of foliation. The foliations are defined by: (a) compositional layering; (b) grain-size variation; (c) closely spaced, approximately parallel discontinuities such as microfaults or fractures; (d) preferred orientation of grain boundaries; and (e) preferred orientation of platy minerals or lenticular mineral aggregates. These various microstructures can be combined and (f) shows a combination (a + e) that is very common in both sedimentary and metamorphic rocks.

Axial Plane Foliation



Divergent Foliation



▪ **Lineation**

Lineation: penetrative linear fabric. We will focus on those that are related to deformation.

How does it differ from other linear structures we have talked about, like slickenlines on a fault surface?

• **Types of lineation:**

- 1) Intersection
- 2) Crenulation
- 3) Mineral
- 4) Stretching

Intersection lineation: Intersection of two planar features- an "apparent" lineation in that there is no fabric that is linear.




Mineral lineation: preferred alignment of minerals due to deformation and/or recrystallization during deformation

Stretching lineation: elongation of minerals due to "stretching" deformation



S-C fabrics- occur in L-S tectonites and serve as excellent sense-of-shear indicators



S-Surfaces- planes of schistosity/foliation (flattening)

C-Surfaces- planes of maximum shear "shear bands"

C comes from cisaillement, French for shear

References

▪ Essential references

Anderson, r. n., et al. (1977), The mechanisms of heat transfer through the floor of the Indian ocean, *J. Geophys. Res.*, *82*, 3391–3409.

Andreas fault zone, J. (1992), *Geophys. Res.*, *85*, 6185–6222.

Davis, G.H.& Reynolds, S.J.D; Structural geology of rocks and regions. 776 pp., John Wiley & Sons, New York, 1996.

Mattauer, M. ; Ce que disent les pierres. Belin. 144 pp., Belin., 1998

Ramsay, J.G. & Hubner, M.; The techniques of modern structural geology. Strain analysis. Vol.1, 307 pp., Academic Press., 1993

Ramsay, J.G. & Hubner, M.; The techniques of modern structural geology. Folds and fractures., Vol.2, 700 pp., 1983. ISBN: Academic Press

▪ Additional references

Brodsky, E. E., and H. Kanamori (2001), Elastohydrodynamic lubrication of faults, *Journal of Geophysical Research-Solid Earth*, *106*, 16374–16305.

Chester, F. M., et al. (2004), Structure of large-displacement Strike-Slip Fault zones in the Brittle Continental Crust, in *Rheology and Deformation of the Lithosphere at Continental Margins*, edited by g. Karner, et al., pp. 223–260, Columbia university Press, New York.

Chester, F. M., et al. (1993), Internal Structure and Weakening Mechanisms of the San-Andreas Fault, *J. Geophys. Res.-Solid Earth*, *98*, 771–786.

Davis, John C., (1986), Statistics and data analysis in geology: 2nd ed., John Wiley & Sons, New

York , 646p.

di Toro, g., and g. Pennacchioni (2005), Fault plane processes and mesoscopic structure of a strong-type seismogenic fault in tonalites (Adamello batholith, Southern Alps), *Tectonophysics*, *402*, 55–80.

Faulkner, d. r., et al. (2003), on the internal structure and mechanics of large strike-slip fault zones: field observations of the Carboneras fault in southeastern Spain, *Tectonophysics*, *367*, 235–251.

Hardebeck, J. I., and E. Hauksson (1999), role of fluids in faulting

inferred from stress field signatures, *Science*, *285*, 236–239.

Greiling, R.O., Abdeen, M.M., Dardir, A.A., El Akhal, H., El Ramly, M.F., Kamal El Din, G.M., Osman, A.F., Rashwan, A.A., Rice, A.H.N., Sadek, M.F., (1994), A structural synthesis of the Proterozoic Arabian-Nubian Shield in Egypt. *Geologie Rundschau*, *83*, 484-501.

Structural Geology Course for 3rd Year Students - Geology Department

Rice, J. r. (1992), Fault stress states, pore pressure distributions, and the weakness of the San Andreas fault, in *Fault Mechanics and Transport Properties of Rocks*, edited by B. Evans and T.-F. Wong, pp. 475–504, Academic Press, London

Scholz, C. H., and T. C. Hanks (2004), The Strength of the San Andreas Fault: A discussion, in *Rheology and Deformation of the Lithosphere at Continental Margins*, edited by g. d. karner, et al., pp. 261–283, Columbia university Press, new york.

Pearce, J.A., (2003). Supra-subduction zone ophiolites: the search for modern analogues. In: Dilek, Y., Newcomb, S. (Eds.), *Ophiolite Concept and the Evolution of Geological Thought*. Geological Society of America Special Paper 373, pp.295-269.

Marshak, Stephen and Mitra, Gautam, (1988), Basic methods of structural geology: Prentice Hall, Englewood Cliffs, New Jersey, 446p.

Ramsay, John G., 1967, Folding and Fracturing of Rocks: New York, McGraw-Hill, 568p.

Stern, R.J., (2005), Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that modern episodes of subduction tectonics began in Neoproterozoic time. *Geology* 33, 557–560.

Stern, R.J., (2008), Neoproterozoic crustal growth: the solid Earth system during a critical episode of Earth history. *Gondwana Research* 14, 33–50.

<http://www.utahpictures.com/Checkerboard.html>.

<http://volcanoes.usgs.gov/Products/Pglossary/PillowLava.html>

<http://volcanoes.usgs.gov/Products/Pglossary/ancientseq.html>

<http://www.geology.washington.edu/~cowan/faultrocks.html#photo>

Scholz, C. H., *The Mechanics of Earthquakes and Faulting*, 2nd. ed., Cambridge University Press, 471 p., 2002

Coney, P. J., Cordilleran metamorphic core complexes; an overview, *in*:Crittenden, M.,D., Jr., Coney, P. J., and G. H. Davis (eds), *Cordilleran metamorphic core complexes*, Geological Society of America Memoir 153, 7-31. 1980.

<http://home.earthlink.net/~rhaughy/ROCKS.HTM>

<http://earth.leeds.ac.uk/assynt/quartzmfr.htm>

<http://www.lpl.arizona.edu/~rlorenz/pseud.html>

<http://www.geolab.unc.edu/Petunia/IgMetAtlas/meta-micro/mylonite.X.html>

<http://www.nps.gov/deva/pphtml/maps.html>

Structural Geology Course for 3rd Year Students - Geology Department

<http://www.geophysics.rice.edu/departement/research/julia1/julia1.html>

<http://earth.leeds.ac.uk/mtb/background/nwmap.htm>

<http://earth.leeds.ac.uk/assynt/quartzmfr.htm>